

The Brain Finger Protein Gene (*ZNF179*), a Member of the RING Finger Family, Maps Within the Smith-Magenis Syndrome Region at 17p11.2

Toshiyuki Kimura,¹ Yoshiki Arakawa,¹ Satoshi Inoue,^{2,6} Yoshimitsu Fukushima,³ Ikuko Kondo,⁴ Kumiko Koyama,⁵ Takayuki Hosoi,⁶ Akira Orimo,² Masami Muramatsu,⁶ Yusuke Nakamura,⁵ Tatsuo Abe,¹ and Johji Inazawa^{1*}

¹Department of Hygiene, Kyoto Prefectural University of Medicine, Kyoto, Japan

²Department of Geriatrics, Faculty of Medicine, University of Tokyo, Tokyo, Japan

³Department of Hygiene and Medical Genetics, Shinshu University School of Medicine, Matsumoto, Japan

⁴Department of Hygiene, Ehime University School of Medicine, Ehime, Japan

⁵Laboratory of Molecular Medicine, Institute of Medical Science, University of Tokyo, Tokyo, Japan

⁶Department of Biochemistry, Saitama Medical School, Saitama, Japan

Smith-Magenis syndrome (SMS) is caused by a microdeletion of 17p11.2 and comprises developmental and growth delay, facial abnormalities, unusual behavior and sleep problems. This phenotype may be due to haploinsufficiency of several contiguous genes. The human brain finger protein gene (*ZNF179*), a member of the RING finger protein family, has been isolated and mapped to 17p11.2. FISH analyses of metaphase or interphase chromosomes of 6 patients with SMS show that *ZNF179* was deleted in one of the 2 homologs (17p11.2), indicating a possible association of the defect of this gene with the pathogenesis of SMS. Furthermore, using a prophase FISH ordering system, we sublocalized *ZNF179* proximally to *LLGL* which lies on the critical region for SMS. Am. J. Med. Genet. 69:320–324, 1997.

© 1997 Wiley-Liss, Inc.

KEY WORDS: Smith-Magenis syndrome; 17p11.2; gene mapping; FISH; *ZNF179*; *LLGL*

INTRODUCTION

A region of 17p is associated with at least 5 distinct syndromes: Miller-Dieker syndrome (MDS), Charcot-Marie-Tooth disease type 1A (CMT1A), hereditary neuropathy with liability to pressure palsies (HNPP), Sjögren-Larsson syndrome (SLS), and Smith-Magenis

syndrome (SMS). The MDS deletions have been mapped to 17p13.3 [Ledbetter et al., 1989]. CMT1A and HNPP are due to the reciprocal DNA duplication/deletion of a 1.5-Mb region of 17p11.2-p12 [Lupski et al., 1991; Raeymaekers et al., 1991; Pentao et al., 1992; Chance et al., 1993, 1994; Wise et al., 1993], which includes *PMP22*, the gene responsible for both disorders. In addition, based on a linkage analysis in SLS families of non-Swedish ethnic origins, SLS has been mapped to a limited region within 17p12 proximal to CMT1A [Rogers et al., 1995]. In contrast, an SMS microdeletion is present at more proximal in 17p11.2.

Patients with SMS show several nonspecific clinical manifestations such as mental retardation, facial abnormalities, and slow growth in addition to more specific manifestations such as unusual behavior and sleep problems [Smith et al., 1986; Greenberg et al., 1991]. Thus, SMS is considered a contiguous gene syndrome. So far 5 genes have been mapped to the SMS critical region: genes coding small RNA U3 (*snRNA U3*) [Chevallard et al., 1993], the human homolog of the *Drosophila melanogaster* flightless-1 (*FLI1*) [Chen et al., 1995], microfibril-associated glycoprotein (*MFAP4*) [Zhao et al., 1995], cytosolic serine hydroxymethyltransferase (*cSHMT*) [Elsa et al., 1995], and of the murine *mg1-1* (*LLGL*) [Koyama et al., 1996].

The human, mouse and rat genes encoding brain finger protein (bfp), a member of RING finger protein family, have been isolated and localized by fluorescence *in situ* hybridization (FISH) [Matsuda et al., 1996]. Of these, the human BFP gene (*ZNF179*) was mapped to 17p11.2 [Matsuda et al., 1996]. The expression of *ZNF179*-mRNA was detected predominantly in the brain, and bfp proteins were also shown in both mouse and rat brain. To define the participation of this gene in SMS, we performed FISH experiments on metaphase and interphase chromosomes from 6 SMS patients whose karyotype analyses had shown microdeletions of 17p11.2. Furthermore, the precise sublocalization of

Contract grant sponsor: The Ministry of Education, Science and Culture, Japan.

*Correspondence to: Johji Inazawa, M.D., Ph.D., Department of Hygiene, Kyoto Prefectural University of Medicine, Kamigyo-ku, Kyoto 602, Japan.

Received 26 April 1996; Accepted 31 July 1996

this gene with respect to *LLGL* was determined by prophase FISH.

SUBJECTS, MATERIALS AND METHODS

SMS Patients

Six unrelated SMS patients examined in this study were described elsewhere [Kondo et al., 1991; Koyama et al., 1996]. All of them show visible microdeletions of sub-band p11.2 on either 17-homolog upon high-resolution banding. Four of these patients (KY, MT, MS and AS) lack *LLGL* in one copy of chromosome 17 [Koyama et al., 1996]. All 6 patients show the unusual behavior typical of SMS; details of the clinical manifestations of the 2 patients (KS and HN) were described by Kondo et al. [1991].

Mapping of ZNF179 to Chromosomes From 6 SMS Patients

The 10-kb lambda phage clone, hgA, contains the sequences for RING finger domain of human brain finger protein [Matsuda et al., 1996]. The references for the positions of 17cen and 17p11.2 in multicolor FISH were *LLGL*, as well as CI17-321 and CI17-536 cosmids which are mapped to 17cen and 17p11.2, respectively [Inazawa et al., 1993]. The cosmids of *LLGL* and CI17-536 are localized within 17p11.2 microdeletions in the 6 patients [Koyama et al., 1996].

Chromosome slides were prepared from EB-virus-transformed lymphoblastoid cell lines from the 6 patients. FISH was carried out according to Inazawa et al. [1993]. In brief, DNA probes of hgA (*ZNF179*) and CI17-321 were labeled with biotin-16-dUTP and digoxigenin-11-dUTP (both from Boehringer-Mannheim, Germany) by nicktranslation, respectively. The labeled probes were denatured at 75°C for 10 min and hybridized to the denatured chromosomes at a final concentration of 25 ng/μl in a mixture of 50% formamide, 10% dextran sulfate (Sigma, St Louis, USA), 2xSSC, sonicated salmon sperm DNA (2 μg/μl), *Escherichia coli* tRNA (2 μg/μl), and Cot-1 DNA (125 ng/μl). The hybridization signals were detected with FITC-avidin and antidigoxigenin rhodamine (both from Boehringer-Mannheim); chromosomes were then counterstained with DAPI (1 μg/ml), and mounted in an anti-fade solution containing 1% DABCO (Sigma). Three-color FISH with hgA, *LLGL* and CI17-536 was performed as in our previous report [Inazawa et al., 1992]. Using hgA double-labeled with biotin-16-dUTP and digoxigenin-11-dUTP, *ZNF179* was visualized as yellow pseudocolor signals generated by FITC and rhodamine. Multicolor signals were simultaneously observed using a Nikon FXA fluorescent microscope equipped with double or triple band-pass filters (Nikon, Tokyo, Japan).

Ordering of ZNF179 and LLGL on 17p11.2 by Prophase FISH

To determine the physical order of *ZNF179* and *LLGL* on sub-band 17p11.2, we used a prophase FISH ordering system [Inazawa et al., 1994]. In brief, 0.5 μg of each *LLGL* cosmid and *ZNF179* lambda phage were labeled with biotin-16-dUTP and digoxigenin-11-

dUTP, respectively. Elongated prophase chromosomes were prepared from cultured lymphocytes by means of thymidine synchronization and bromodeoxyuridine release in the presence of topoisomerase II inhibitor ICRF 154 [Inazawa et al., 1994]. Hybridization was performed at 37°C overnight on denatured-chromosome slides. After washing, the slides were incubated at 37°C with FITC-avidin and anti-digoxigenin rhodamine. The slides were washed at room temperature for 10 min each in 4xSSC, 4xSSC containing 0.05% Triton X-100, and 4xSSC, then counterstained with DAPI (1 μg/ml). To determine the order of *ZNF179* and *LLGL* probes by multi-color FISH, FITC and rhodamine signals were simultaneously detected through a triple band-pass filter, then the contours on DAPI-stained prophase chromosomes were delineated through a UV-2A filter (Nikon). The order and orientation of signals on sub-band 17p11.2 were determined by pairwise comparison along the long axis of chromatids.

RESULTS

FISH of ZNF179 in 6 SMS Patients

More than 80 cells (over 30 metaphases and 50 interphases) were examined on each of the 6 patients. FITC signals specific for *ZNF179* were detected on only one copy of chromosome 17 of metaphase cells of all 6 patients, although rhodamine red signals specific for the centromere of chromosome 17 were detected on both homologs (Fig. 1a). In addition, three-color FISH with *ZNF179*, *LLGL* and CI17-536 detected only one spot of each at one of the 2 homologs in interphase nuclei from the 6 patients (Fig. 1b). These results indicated that one copy of *ZNF179* together with *LLGL* is lacking in these patients.

Ordering of ZNF179 and LLGL by Prophase FISH

To determine the physical order of *ZNF179* and *LLGL*, we examined prophase chromosome 17 showing two different colored signals of a pair of *ZNF179* and *LLGL* cosmid probes. Relative positions of the signals were scored along the long axis of chromosomes. Of 20 prophase chromosomes examined, 17 had 17pter-green (FITC)-red (rhodamine)-17cen signals on both chromatids (Fig. 2) and FITC overlapped rhodamine signals on 17p11.2 in three. These results indicate that the physical order of *ZNF179* and *LLGL* is 17pter-*LLGL*-*ZNF179*-17cen.

DISCUSSION

We have shown that the human brain finger protein gene (*ZNF179*), a member of RING finger protein family, was deleted on one copy of chromosome 17 in all of the 6 SMS patients. SMS patients with an interstitial microdeletion of 17p11.2 have peripheral neuropathy, seizure, sleep disturbance, and behavior abnormalities including hyperactivity and self-mutilation [Smith et al., 1986; Greenberg et al., 1991], in addition to non-specific mental retardation, facial anomalies and developmental delay. These phenotypes suggest that the SMS critical region contains important gene(s) associ-




Fig. 1. FISH analysis in SMS patients. **a:** (Left) An FITC signal (arrow) specific for *ZNF179* was detected on one of two chromosome 17 homologs in an SMS patient (AS), although rhodamine red signals (open arrow) specific for 17cen were detected on both homologs. Right: The same metaphase chromosomes were counterstained with DAPI. Arrowhead indicates chromosomes 17. **b:** Representative nucleus from an SMS patient (HN) showing three-color FISH with probes of hgA (*ZNF179*), *LLGL* and CI17-536. Only one spot of each probe was detected on one chromosome 17p11.2, indicating that *ZNF179* (yellow) and *LLGL* (red) together with CI17-536 (green) are missing from one copy of chromosome 17 in this patient.




Fig. 2. Ordering of *ZNF179* and *LLGL* by prophase FISH. **Right:** A pair of FITC-labeled hgA (*ZNF179*) and rhodamine-labeled *LLGL* cosmid were detected in the order: 17pter-*LLGL*-*ZNF179*-17cen. **Left:** The same chromosomes were counterstained with DAPI.

ated with neuronal development, differentiation and function.

The brain finger protein (bfp) of the rat was the first to be isolated and characterized. The expression of bfp mRNA was predominantly detected in the brain, and bfp proteins were also shown in the rat and mouse brain. The bfp protein was not detected in untreated P19 embryonic carcinoma cells but was evident after exposure to retinoic acid when the cells differentiated into a neural cell-like form. These findings suggest that the bfp protein is up-regulated during the neural differentiation of P19 cells by retinoic acid and that it is a putative transcription regulator in the brain. Hence, human BFP may also be important in neuronal development and differentiation, and the haploinsufficiency of this gene might be involved in the pathogenesis of SMS.

Since a cytogenetically detectable deletion encompasses ≥ 2 –3 Mb [Ledbetter and Cavenee, 1989], it is likely that several genes lie on the SMS deleted region. Extensive studies of physical mapping spanning the SMS deleted region have been reported [Greenberg et al., 1991; Guzzetta, et al., 1992; Chevillard et al., 1993]. Chevillard et al. [1993] suggested that the SMS critical region is over 2–2.5 Mb, and variations of the sizes of microdeletions are likely to be due to the differences of the proximal breakpoints. Our study of prophase FISH showed that *ZNF179* is proximal to *LLGL* at 17p11.2. SMS shows phenotypic variations similar to those of other congenital chromosome abnormalities [Smith et al., 1986; Kondo et al., 1991; Greenberg et al., 1991; Zori et al., 1993; Finucane et al., 1993, 1994; Colley et al., 1993]. Thus, the presence or absence of *ZNF179* within the critical region for SMS should be examined in more SMS patients, and studies of biological function of this gene are required to determine if *ZNF179* is involved in eliciting the SMS-specific phenotypes.

ACKNOWLEDGMENTS

We appreciate useful discussions with Drs. Yoichi Matsuda, Tada-aki Hori, and Hirofumi Ohashi. This work was supported in part by a Grant-in-Aid from the Ministry of Education, Science and Culture, Japan.

REFERENCES

- Chance PF, Alderson MK, Leppig KA, Lensch MW, Matsunami N, Smith B, Swanson PD, Odelberg SJ, Distech CM, Bird TD (1993): DNA deletion associated with hereditary neuropathy with liability to pressure palsies. *Cell* 72:143–151.
- Chance PF, Abbas N, Lensch MW, Pentao L, Roa BB, Patel PI, Lupski JR (1994): Two autosomal dominant neuropathies result from reciprocal DNA duplication/deletion of a region on chromosome, 17. *Hum Molec Genet* 3:223–228.
- Chen K-S, Gunaratne PH, Hoheisel JD, Young IG, Miklos GLG, Greenberg F, Shaffer LG, Campbell HD, Lupski JR (1995): The human homologue of the *Drosophila melanogaster* flightless-1 gene (*fli1*) maps within the Smith-Magenis microdeletion critical region in 17p11.2. *Am J Hum Genet* 56:175–182.
- Chevillard C, Paslier DL, Passage E, Ougen P, Billaut A, Boyer S, Mazan S, Bachelier JP, Vingal A, Cohen D, Fontes M (1993): Relationship between Charcot-Marie-Tooth 1A and Smith-Magenis regions: *snU3* may be a candidate gene for the Smith-Magenis syndrome. *Hum Molec Genet* 2:1235–1243.
- Colley AF, Leversha MA, Voullaire LE, Rogers JG (1990): Five cases demonstrating the distinctive behavioral features of chromosome deletion 17 (p11.2p11.2) (Smith-Magenis syndrome). *J Pediatr Child Health* 26:17–21.
- Elsea SH, Juyal RC, Jiralerspong S, Finucane BM, Pandolfo M, Greenberg F, Baldini A, Stover P, Patel PI (1995): Haploinsufficiency of cytosolic serine hydroxymethyltransferase in the Smith-Magenis syndrome. *Am J Hum Genet* 57:1342–1350.
- Finucane BM, Jaeger ER, Kurtz MB, Weinstein M, Scott CI (1993): Eye abnormalities in the Smith-Magenis contiguous gene deletion syndrome. *Am J Med Genet* 45:443–446.
- Finucane BM, Konar D, Haas-Givler, Kurtz MB, Scott CI (1994): The spasmodic upper-body squeeze: a characteristic behavior in Smith-Magenis syndrome. *Dev Med Child Neurol* 36:70–83.
- Greenberg F, Guzzetta V, Montes de Oca-Luna R, Magenis RE, Smith ACM, Richter SF, Kondo I, Dobyns WB, Patel PI, Lupski JR (1991): Molecular analysis of the Smith-Magenis syndrome: a possible contiguous gene syndrome associated with del(17)(p11.2). *Am J Hum Genet* 49:1207–1218.
- Guzzetta V, Franco B, Trask BJ, Zhang H, Saucedo-Cardenas O, Montes de Oca-Luna R, Greenberg F, Chinault AC, Lupski JR, Patel PI (1992): Somatic cell hybrids, sequence-tagged sites, simple repeat polymorphisms, and yeast artificial chromosomes for physical and genetic mapping of proximal 17p. *Genomics* 13:551–559.
- Inazawa J, Ariyama T, Abe T (1992): Physical ordering of three polymorphic DNA markers spanning the regions containing a tumor suppressor gene of renal cell carcinoma by three-color fluorescent in situ hybridization. *Jpn J Cancer Res* 83:1248–1252.
- Inazawa J, Saito H, Ariyama T, Abe T, Nakamura Y (1993): High-resolution cytogenetic mapping of 342 new cosmid markers including 43 RFLP markers on human chromosome 17 by fluorescence in situ hybridization. *Genomics* 17:153–162.
- Inazawa J, Ariyama T, Tokino T, Tanigami A, Nakamura Y, Abe T (1994): High resolution ordering of DNA markers by multi-color fluorescent in situ hybridization of prophase chromosomes. *Cytogenet Cell Genet* 65:130–135.
- Kondo I, Matsuura S, Kuwajima K, Tokashiki M, Izumikawa Y, Naritomi K, Niikawa N, Kajii T (1991): Diagnostic hand anomalies in Smith-Magenis syndrome: four new patients with del(17)(p11.2p11.2). *Am J Med Genet* 41:225–229.
- Koyama K, Fukushima Y., Inazawa J., Tomotsune D., Takahashi N., Nakamura Y (1996): The human homologue of the murine *Llgh* gene (*LLGL*) maps within the Smith-Magenis syndrome region in 17p11.2. *Cytogenet Cell Genet* 72:78–82.
- Ledbetter DH, Cavenee WK (1989): Molecular Cytogenetics: interface of cytogenetics and monogenic disorders. In Scriver CR, Beaudet AL, Sly WS, Valle D (eds) "The Metabolic Basis of Inherited Disease," 6th ed. New York: McGraw Hill, pp 343–371.
- Ledbetter DH, Ledbetter SA, van Tuinen P, Summers KM, Robinson TJ, Nakamura Y, Wolff R, White R, Barker DF, Wallace MR, Collins FS, Dobyns WB (1989): Molecular dissection of a contiguous gene syndrome: frequent submicroscopic deletions, evolutionary conserved sequences, and a hypomethylated "island" in the Miller-Dicker chromosome region. *Proc Natl Acad Sci USA* 86:5136–5140.
- Lupski JR, Montes de Oca-Luna R, Slaugenhaupt S, Pentao L, Guzzetta V, Trask BJ, Saucedo-Cardenas O, Barker DF, O'Connell P, Chance PF (1991): DNA duplication associated with Charcot-Marie-Tooth disease type 1A. *Cell* 66:219–232.
- Matsuda Y, Inoue S, Seki N, Hosoi T, Orimo A, Muramatsu M, Hori T (1996): Chromosome mapping of human (*ZNF179*), mouse and rat genes for brain finger protein (bfp), a member of the RING finger family. *Genomics* 33:325–327.
- Pentao L, Wise CA, Chinault AC, Patel PI, Lupski JR (1992): Charcot-Marie-Tooth type 1A duplication appears to arise from recombination at repeat sequences flanking the 1.5 Mb monomer unit. *Nature Genet* 2:292–300.
- Raeymaekers P, Timmerman V, Nelis E, De Jonghe P, Hoogendijk JE, Baas F, Barker DF, Martin JJ, De Visser M, Bolhuis PA, Van Broeckhoven C, the HMSN Collaborative Research Group (1991): Duplication in chromosome 17p11.2 in Charcot-Marie-Tooth neuropathy type 1a (CMT1a). *Neuromusc Disord* 1:93–97.
- Rogers G R, Rizzo W B, Zlotogorski A, Hashem N, Lee M, Compton J G, Bale SJ (1995): Genetic homozygosity in Sjögren-Larsson syndrome: Linkage to chromosome 17p in families of Different non-Swedish ethnic origins. *Am J Hum Genet* 57:1123–1129.
- Smith ACM, McGavran L, Robinson J, Waldstein G, Macfarlane J, Zonona

- J, Reiss J, Lahr M, Allen L, Magenis E (1986): Interstitial deletion of (17)(p11.2p11.2): report of six additional patients with a new chromosome deletion syndrome. *Am J Med Genet* 24:421–432.
- Wise CA, Garcia CA, Davis SN, Heju Z, Pentao L, Patel PI, Lupski JR (1993): Molecular analysis of unrelated Charcot-Marie-Tooth (CMT) disease patients suggest a high frequency of the CMT1A duplication. *Am J Hum Genet* 53:853–863.
- Zhao Z, Lee C-C, Jiralerspong S, Juyal RC, Lu F, Baldini A, Greenberg F, Caskey CT, Patel PI (1995): The gene for a human microfibril-associated glycoprotein is commonly deleted in Smith-Magenis syndrome patients. *Hum Molec Genet* 4:4:589–597.
- Zori RT, Lupski JR, Heju Z, Greenberg F, Killian JM, Gray BA, Driscoll DJ, Patel PI, Zackowski JL (1993): Clinical, cytogenetic and molecular evidence for an infant with Smith-Magenis syndrome born from a mother having a mosaic 17p11.2p12 deletion. *Am J Med Genet* 47:504–511.